

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 September 2005 (29.09.2005)

PCT

(10) International Publication Number
WO 2005/090286 A1

(51) International Patent Classification⁷: **C07C 235/44**,
235/46, A61K 31/166, A61P 3/04

(21) International Application Number:
PCT/US2005/007051

(22) International Filing Date: 8 March 2005 (08.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04380058.0 12 March 2004 (12.03.2004) EP
60/553,184 15 March 2004 (15.03.2004) US

(71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DE LA TORRE, Marta, Garcia** [ES/ES]; Lilly, S.A., Avenida de la Industria 30, 28108 Alcobendas-Madrid (ES). **MITCH, Charles, Howard** [US/US]; 3210 Grove Parkway, Columbus, IN 47203 (US).

(74) Agents: **GINAH, Francis, O.** et al.; Eli Lilly and Company, P.O. Box 6288, Indianapolis, IN 46206-6288 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

[Continued on next page]

(54) Title: OPIOID RECEPTOR ANTAGONISTS



(57) Abstract: A compound of the formula (I) wherein the variables X1, X2, R1 to R7 including R3', j, k, p, y, and z, are as defined or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixtures thereof, useful for the treatment, prevention or amelioration of obesity and Related Diseases is disclosed.

**Published:**

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

OPIOID RECEPTOR ANTAGONISTS

The present invention is in the field of medicinal chemistry. The invention relates specifically to compounds useful as opioid antagonists, methods of treatment, methods of using, and pharmaceutical compositions thereof.


Background

Three types of opioid receptors, mu, kappa, and delta opioid receptors are generally reported. Recent evidence points to the interactions between receptor dimer combinations of mu, kappa and/or delta receptors (called heterodimers) as also contributing to opioid activity. Opioid receptors and their normal regulation or lack thereof, has been implicated in disease states including irritable bowel syndrome, nausea, vomiting, pruritic dermatoses, depression, smoking and alcohol addiction, sexual dysfunction, stroke and trauma in animals. Therefore it is not surprising that the ability to antagonistically bind opioid receptors has been shown to produce ameliorative, preventative and/or treatment effects in animals including humans afflicted with one or more of these disease states.


More recently, certain antagonists of the opioid receptors have been found to increase metabolic energy consumption, and reduction of weight in obese rats while maintaining muscle mass. These findings indicate that an effective opioid antagonist may be useful in preventing, treating and/or ameliorating the effect of obesity. Considering the percentage of the population that is obese in Western societies and the indirect costs associated with treating the effects and symptoms of obesity and Related Diseases, the importance of these findings cannot be overstated.

Though many opioid antagonists have been disclosed, the search continues for alternative and/or improved or more effective antagonists having an overall benefit to the patient with little or no major side effects. U.S. Patent No. 4,891,379 disclosed phenylpiperidine opioid antagonists useful for the treatment of diabetes and obesity. In particular, U.S. patent 4,891,379 disclosed the compound LY 255582 represented by the structure

2



U.S. Patent No. 4,191,771 also disclosed compounds useful as opioid antagonists. Also, bicyclic analogs of phenyl piperidine have been prepared and reported as opioid antagonists in Wentland, et al., *Biorganic and Medicinal Chemistry Letters* 11 (2001) 623-626; see also Wentland, et al., *Bioorganic and Medicinal Chemistry Letters* 11 (2001) 1717-1721. Finally, European Patent application number EP 1 072592A2 filed May 18, 2000, discloses phenylpiperidine compounds of formula 1




1

wherein A, D, R¹, R², R³, X, and n have meanings given in the description, which are useful in the prophylaxis and in the treatment of diseases mediated by opioid receptors such as pruritus.


U.S patent No. 6,140,352 discloses the compound of formula

Formula 1



wherein the variables X_1 , X_2 , X_3 , R_1 , R_3 , R_4 , R_5 and R_6 are as described therein, as agonists of the beta adrenergic receptor useful for the treatment of diabetes and obesity.

PCT application WO 9215304 discloses the compounds of formula I




Compound 1 above, encompasses azacyclic and heterocyclic compounds for treatment of cerebral ischemia.

Regardless of these and other disclosures of compounds useful as opioid receptor antagonists, or useful for the treatment of obesity, and/or diabetes by other mechanisms, or having structures partially close to the compounds of the present invention there remains an unmet medical need for useful, safe, effective and/or alternate treatments or prophylaxis of diseases associated with opioid receptors, particularly obesity and Related Diseases.

Summary of the Invention

The present invention provides a compound of the formula (I)



p is 0, 1, or 2;

y is 0, 1, or 2; and z is 0, 1, or 2;

j and k are each independently 0, 1, or 2 provided that j and k are not simultaneously equal to 0;

X_1 and X_2 are each independently is CH, or N;

R^1 and R^2 are independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl, C_1 - C_{10} alkylaryl, SO_2R^8 , $(CH_2)_nC(O)NR^8R^8$, SO_2C_1 - C_{10} alkylaryl,

$\text{SO}_2\text{C}_1\text{-C}_8$ alkylheterocyclic, $\text{C}_4\text{-C}_{10}$ alkylcycloalkane, $(\text{CH}_2)_n\text{C}(\text{O})\text{OR}^8$, and $(\text{CH}_2)_n\text{C}(\text{O})\text{R}^8$; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to two groups independently selected from $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, phenyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $\text{C}_1\text{-C}_8$ alkylaryl, and $\text{C}(\text{O})\text{C}_1\text{-C}_8$ alkyl; and wherein R^1 and R^2 may optionally combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may further have substituents selected from the group consisting of oxo, amino, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, $\text{C}_2\text{-C}_8$ alkynyl, phenyl, $\text{C}_1\text{-C}_3$ alkylaryl, $\text{C}(\text{O})\text{C}_1\text{-C}_8$ alkyl, $\text{CO}(\text{O})\text{C}_1\text{-C}_8$ alkyl, halo, $\text{C}_1\text{-C}_3$ haloalkyl;

R^3 and $\text{R}^{3'}$ are each independently selected from hydrogen, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, $\text{C}_2\text{-C}_8$ alkynyl, phenyl, aryl, $\text{C}_1\text{-C}_8$ alkylcycloalkyl, and $\text{C}_1\text{-C}_8$ alkylaryl;

R^4 and R^5 are each independently selected from hydrogen, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, $\text{C}_2\text{-C}_8$ alkynyl, $\text{C}_1\text{-C}_8$ alkoxy, halo, $\text{C}_1\text{-C}_8$ haloalkyl, phenyl, aryl, $\text{C}_1\text{-C}_8$ alkylaryl, $(\text{CH}_2)_m\text{NSO}_2\text{C}_1\text{-C}_8$ alkyl, $(\text{CH}_2)_m\text{NSO}_2$ phenyl, $(\text{CH}_2)_m\text{NSO}_2$ aryl, $-\text{C}(\text{O})\text{C}_1\text{-C}_8$ alkyl, and $-\text{C}(\text{O})\text{OC}_1\text{-C}_8$ alkyl; wherein each R^4 and R^5 is attached to its respective ring only at carbon atoms; wherein m is 1 or 2; and n is 1, 2, or 3;

R^6 and R^7 are each independently selected from hydrogen, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, $\text{C}_2\text{-C}_8$ alkynyl, $\text{C}(\text{O})\text{C}_1\text{-C}_8$ alkyl, $\text{SO}_2\text{C}_1\text{-C}_8$ alkyl, $\text{SO}_2\text{C}_1\text{-C}_8$ alkylaryl, $\text{SO}_2\text{C}_1\text{-C}_8$ alkylheterocyclic, aryl, $\text{C}_1\text{-C}_8$ alkylaryl, $\text{C}_3\text{-C}_7$ cycloalkane, $\text{C}_1\text{-C}_6$ alkylcycloalkane, $(\text{CH}_2)_m\text{C}(\text{O})\text{OR}^8$, $(\text{CH}_2)_m\text{C}(\text{O})\text{R}^8$, $(\text{CH}_2)_m\text{C}(\text{O})\text{NR}^8\text{R}^8$, and $(\text{CH}_2)_m\text{NSO}_2\text{R}^8$; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to two groups independently selected from $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, phenyl, and $\text{C}_1\text{-C}_8$ alkylaryl; and wherein R^6 and R^7 may independently combine with each other, and with the nitrogen atom to which they are attached to form a 4, 5, 6, or 7-membered nitrogen containing heterocycle which nitrogen containing heterocycle may optionally have substituents selected from the group consisting of oxo, amino, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, $\text{C}_2\text{-C}_8$ alkynyl, phenyl, and $\text{C}_1\text{-C}_8$ alkylaryl;

R^8 is independently selected from hydrogen, $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_2\text{-C}_8$ alkenyl, phenyl, benzyl, and $\text{C}_5\text{-C}_8$ alkylaryl; or a pharmaceutically acceptable salt, solvate, prodrug, enantiomer, racemate, diastereomer, or mixture of diastereomers thereof.

The present invention also provides a method for the prevention, treatment and/or amelioration of the symptoms of obesity and Related Diseases comprising administering

a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

The present invention also provides a pharmaceutical formulation comprising a compound of formula I in association with a carrier, diluent and/or excipient.

The present invention also relates to a method for the treatment and/or prophylaxis of obesity and Related Diseases including eating disorders (bulimia, anorexia nervosa, etc.), diabetes, diabetic complications, diabetic retinopathy, sexual/reproductive disorders, depression, anxiety, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure, sleeping disorders, atherosclerosis, rheumatoid arthritis, stroke, hyperlipidemia, hypertriglycemia, hyperglycemia, hyperlipoproteinemia, substance abuse, drug overdose, compulsive behavior disorders (such as paw licking in dog), and addictive behaviors such as for example, gambling, and alcoholism, comprising administering a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

The present invention relates to a compound of formula (I) useful for the manufacture of a medicament for the treatment, prevention and/or amelioration of symptoms associated with obesity and Related Diseases.

In another embodiment, the present invention relates to a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture thereof, useful as an appetite suppressant.

The present invention relates to a method of achieving weight loss while maintaining lean muscle mass or minimizing the loss of lean muscle mass comprising administering a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture thereof, to a patient in need thereof.

The present invention provides a compound of formula I useful singly or in combination with other agents approved for the treatment, prevention and/or amelioration of obesity and related diseases and symptoms thereof.

Detailed Description of the Invention

As used herein, the term "patient" includes human and non-human animals such as companion animals (dogs and cats and the like) and livestock animals.

The preferred patient of treatment, amelioration and/or prevention of obesity and Related Diseases is a human.

The terms "treating" and "treat", as used herein, include their generally accepted meanings e.g. preventing, prohibiting, restraining, alleviating, ameliorating, slowing, stopping, or reversing the progression or severity of a pathological condition, or sequela thereof, described herein.

The terms "ameliorating" "preventing", "prevention of", "prophylaxis", "prophylactic" and "prevent" are used herein interchangeably and refer to reducing the severity of obesity and Related Diseases and the symptoms associated therewith, in a patient afflicted with same or reducing the likelihood that the recipient of a compound of formula I will be afflicted with or develop any of the pathological conditions or sequela thereof described herein.

As used herein, the term "effective amount" is synonymous with "effective dose" and means an amount of a compound of formula I that is sufficient in one or more administrations for preventing, ameliorating or treating a condition, or detrimental effects thereof, herein described, or an amount of a compound of formula I that is sufficient for antagonizing the opioid receptors to achieve the objectives of the invention.

The term "pharmaceutically acceptable" is used herein as an adjective and means substantially non-deleterious to the recipient patient.

The term "Active Ingredient" as used herein means a compound of formula I or a combination of a compounds of formula I or a combination of a compound of formula I and a co-antagonist of the opioid receptor or a combination a compound of formula I and other effective anti-obesity, weight loss or antidiabetic agent.

The term "formulation", as in pharmaceutical formulation, or "pharmaceutical composition" is intended to encompass a product comprising the Active Ingredient (as defined supra), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients.

Accordingly, the pharmaceutical formulations of the present invention encompass any

effective composition made by admixing a compound of the present invention and a pharmaceutical carrier. The pharmaceutical formulations of the present invention also encompass a compound of the formula I and a pharmaceutically acceptable co-antagonist of opioid receptors useful for the treatment and/or prevention of obesity or Related Diseases.

The term "Related Diseases" as used herein refers to such symptoms, diseases or conditions caused by, exacerbated by, induced by or adjunct to the condition of being obese. Such diseases, conditions and/or symptoms include but are not limited to eating disorders (bulimia, anorexia nervosa, etc.), diabetes, diabetic complications, diabetic retinopathy, sexual/reproductive disorders, depression (particularly that induced by the awareness and loss of self esteem associated with obesity), anxiety, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure, sleeping disorders, atherosclerosis, rheumatoid arthritis, stroke, hyperlipidemia, hypertriglycemia, hyperglycemia, and hyperlipoproteinemia.

As used herein "other agents" approved for the treatment of obesity and /or related disease, or useful for weight loss and/or appetite suppression include but are not limited to Xenical®, Meridia®, Lipitor®, Crestor®, Pravachol®, Zetia®, cannabinoid receptor antagonists, and other opioid receptor antagonists.

The term "suitable solvent" refers to any solvent, or mixture of solvents, inert to the ongoing reaction that sufficiently solubilizes the reactants to afford a medium within which to effect the desired reaction.

The term "mutual solvent" means a solvent that is used to dissolve sufficiently, two or more components of a reaction or mixture separately prior to reaction or mixing, that is a solvent common to more than one reagents or components of a mixture.

The term "nitrogen containing heterocycle" refers to a monocycle which is a 4, 5, 6, or 7-member ring containing 1, 2 or 3 nitrogen atoms in addition to the carbon atoms completing the ring size, or a combination of 1 nitrogen atom and 1, or 2 atoms selected from oxygen, and sulfur in addition to the appropriate number of carbon atoms completing the ring size. A nitrogen containing heterocycle as used here may have 0, 1, 2 or 3 double bonds. A nitrogen containing heterocycle may be attached to or fused to an existing ring substituent thus forming a bicyclic or tricyclic ring system. Nonetheless, the

direct result of the formation of a nitrogen containing heterocycle by the joining of two groups and the nitrogen atom to which they are attached is to form a monocycle.

The term "C₁-C₈ alkyl" or C₁₋₈ alkyl" refers to and includes all groups, structural isomers and /or homologues of alkyl groups having from 1 to 8 carbon atoms. When the term C₁-C₈ alkyl precedes or prefixes another group, the term C₁-C₈ alkyl, only limits the number of carbon atoms in the alkyl component. For example C₁-C₈ alkylaryl means an aryl group having a C₁-C₈ alkyl group substituent such that the number of carbon atoms in the group C₁-C₈ alkylaryl is effectively the number of carbon atoms in the aryl group plus the number of carbon atoms in the C₁-C₈ alkyl group. Similarly, the term "C₁-C₈ alkylcycloalkyl" refers to a cycloalkane group having a C₁-C₈ alkyl substituent, and wherein the entire group C₁-C₈ alkylcycloalkane may itself be a substituent attached at either the alkyl group or the cycloalkyl group to a substrate. The definition and usage applies equally to other homologues of C₁-C₈ such as for example, C₁-C₇, C₁-C₆ etc.

The term "cycloalkane" or "cycloalkyl" means cycloalkanes having from 3 to 8 carbon atoms i.e. from cyclopropane to cyclooctane.

The term "hal" or "halo" as used herein refers to a halogen including fluorine, chlorine, bromine or iodine.

As used herein the terms "alkenyl" refers to straight or branched carbon atoms having 1 or 2 carbon-carbon double bonds.

As used herein the terms "alkynyl" refers to straight or branched carbon atoms having 1 or 2 carbon-carbon triple bonds.

As used herein the term "alkoxy" refers to the group "O-alkyl" wherein alkyl is as defined previously.

The term "aryl" as used herein refers to compounds or groups having the Huckel 4n+2 pi electron arrangement and includes phenyl, benzyl, naphthyl, but excludes carbazoles and other fused tricyclic ring structures.

It is understood by one of skill in the art that where a substituent is absent, a hydrogen atom is indicated to achieve the required valency unless otherwise indicated. For example, if y is 0, then R⁴ is absent, and all applicable positions on the ring have hydrogen atoms to achieve the required valency for atoms in the ring.

As used herein, the term "protecting group" refers to a groups useful for masking reactive sites in a molecule to enhance the reactivity of another group or allow reaction at

another desired site or sites following which the protecting group may be removed. Protecting groups are usually used to protect or mask groups including but not limited to -OH, -NH, and -COOH. Suitable protecting groups are known to one of skill in the art and are described in *Protecting groups in Organic Synthesis*, 3rd edition, Greene, T. W.; Wuts, P.G.M. Eds.; John Wiley and Sons, New York, 1999.

As used herein, the term "solvate" is a form of the compound of the invention wherein a crystal or crystals of a compound of the invention have been formed from a stoichiometric or non-stoichiometric amount of the compound of formula I and a solvent. Typical solvating solvents include for example, water, methanol, ethanol, acetone and dimethylformamide.

In those instances where a compound of the invention possesses acidic or basic functional groups, various salts may be formed which are more water soluble and/or more physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts, include but are not limited to, the alkali and alkaline earth salts such as lithium, sodium, potassium, calcium, magnesium, aluminum and the like. Salts are conveniently prepared from the free acid by treating the acid in solution with a base or by exposing the acid to an ion-exchange resin.

Included within the definition of pharmaceutically acceptable salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention, for example, ammonium, quaternary ammonium, and amine cations, derived from nitrogenous bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, *et al.*, "Pharmaceutical Salts," *J. Phar. Sci.*, 66: 1-19 (1977)). Moreover, the basic group(s) of the compound of the invention may be reacted with suitable organic or inorganic acids to form salts such as acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, hydrobromide, camsylate, carbonate, clavulanate, citrate, chloride, edetate, edisylate, estolate, esylate, fluoride, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrochloride, hydroxynaphthoate, hydroiodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, sulfite, sulfate, mucate, napsylate, nitrate, oleate, oxalate, palmitate, pantothenate, phosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, tosylate, trifluoroacetate, trifluoromethane sulfonate, and valerate.

A compound of the invention as illustrated by formula I may occur as any one of its positional isomers, stereochemical isomers or regio-isomers, all of which are objects of the invention. Certain compounds of the invention may possess one or more chiral centers, and thus, may exist in optically active forms. Likewise, when the compounds contain an alkenyl or alkenylene group, there exist the possibility of cis- and trans-isomeric forms of the compounds. The R- and S- isomers and mixtures thereof, including racemic mixtures as well as mixtures of enantiomers or cis- and trans- isomers, are contemplated by this invention. Additional asymmetric carbon atoms can be present in a substituent group such as an alkyl group. All such isomers as well as the mixtures thereof are intended to be included in the invention. If a particular stereoisomer is desired, it can be prepared by methods well known in the art by using stereospecific reactions with starting materials which contain the asymmetric centers and are already resolved or, alternatively by methods which lead to mixtures of the stereoisomers and subsequent resolution by known methods. For example, a racemic mixture may be reacted with a single enantiomer of some other compound i.e. a chiral resolving agent. This changes the racemic form into a mixture of stereoisomers and diastereomers, because they have different melting points, different boiling points, and different solubilities and can be separated by conventional means, such as crystallization.


The compounds of the present invention have shown inhibition of orexigenic effects, and are thus useful as appetite suppressants either as a single therapy or in conjunction with exercise and other effective appetite suppressing or weight loss medications.

Preferred Embodiments of the Invention

A compound of formula I preferably exists as the free base or a pharmaceutically acceptable salt. More preferred is the hydrochloride salt, the bisulfate salt, mesylate or the oxalic acid salt of the compound of formula I.


For the groups R^1 and R^2

Preferred R^1 and R^2 groups are independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, pentyl, and isopropyl. Also preferred are R^1 and R^2 groups independently selected from the group consisting of methyl, ethyl, propyl, isopropyl, phenyl,



each of which is optionally substituted with a group selected from the group consisting of halogen, C₁-C₈ alkyl, C₁-C₈ haloalkyl, C₁-C₈ thioalkyl, C₁-C₈ alkylamino, phenyl, C₁-C₈ alkylsubstituted phenyl, C₄-C₈ heterocycle or C₁-C₄ alkyl heterocycle; or combine with a group selected from C₁-C₈ alkyl, halogen, C₁-C₈ haloalkyl, C₁-C₈ thioalkyl, C₁-C₈ alkylamino, phenyl, C₁-C₈ alkylsubstituted phenyl, C₄-C₈ heterocycle or C₁-C₄ alkyl heterocycle to form a substituted or unsubstituted bicycle or tricycle.

Also preferred are R¹ and R² groups that combine with each other to form a group selected from the group consisting of



each of which is optionally substituted with a group selected from the group consisting of halogen, C₁-C₈ alkyl, C₁-C₈ haloalkyl, C₁-C₈ thioalkyl, C₁-C₈ alkylamino, phenyl, C₁-C₈ alkylsubstituted phenyl, C₄-C₈ heterocycle or C₁-C₄ alkylheterocycle.

Preferred R³ and R^{3'} Groups

A preferred R^3 is hydrogen. A preferred $R^{3'}$ group is selected from hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, phenyl and benzyl. More preferably, both R^3 and $R^{3'}$ are hydrogen.

Preferred R^4 Groups

A preferred R^4 group is selected from the group consisting of hydrogen, halo, C_1 - C_5 alkyl, C_1 - C_5 haloalkyl, C_1 - C_5 alkoxy, C_1 - C_5 alkylamino, phenyl, C_1 - C_5 alkylphenyl, C_1 - C_5 alkylcycloalkyl, and C_1 - C_5 thioalkyl. More preferred is a R^4 group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, chloro, fluoro, trifluoromethyl, methoxy, ethoxy, thiomethyl, phenyl, and benzyl. Most preferred is an R^4 group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, fluoro, chloro, trifluoromethyl, methoxy, ethoxy, propoxy, isopropoxy, and benzyl.

Though the groups R^4 and a R^5 may exist as multiple substituents on their respective ring substrates, a preferred embodiment of the invention involves compounds wherein each of R^4 , and R^5 are independently absent, or singly substituted on their respective ring substrates.

Preferred R^5 Groups

A preferred R^5 group is selected from the group consisting of hydrogen, halo, C_1 - C_5 alkyl, C_1 - C_5 haloalkyl, C_1 - C_5 alkoxy, C_1 - C_5 alkylamino, phenyl, C_1 - C_5 alkylphenyl, C_1 - C_5 alkylcycloalkyl, and C_1 - C_5 thioalkyl. More preferred is an R^5 group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, chloro, fluoro, trifluoromethyl, methoxy, ethoxy, thiomethyl, phenyl, and benzyl. A most preferred R^5 group is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, fluoro, chloro, trifluoromethyl, methoxy, ethoxy, trifluoromethoxy, and benzyl.

Preferred R^6 and R^7 Groups

Preferred are R^6 and R^7 groups independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, pentyl, isopropyl, phenyl and benzyl.

Also preferred are compounds of formula I wherein R^6 and R^7 independently combine with each other, and with the nitrogen atom to which they are attached to form a 4, 5, 6, or 7-membered nitrogen containing heterocycle which nitrogen containing

heterocycle optionally has substituents selected from the group consisting of oxo, amino, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, C₁-C₈ alkylaryl, C(O)C₁-C₈ alkyl, CO(O)C₁-C₈ alkyl, hydroxy, C₁-C₈ alkoxy, halo, and haloalkyl. Most preferred are compounds of the invention wherein R⁶ and R⁷ are both hydrogen.

Preferred values for n, m, and p, j, k, y, z

A preferred value for n is 0, 1 or 2.

A preferred value for m is 1 or 2.

A preferred value for p is 0, 1, or 2. More preferred is p = 1.

A preferred value for j is 1

A preferred value for k is 1

A preferred value for y is 0, or 1

A preferred value for z is 0, or 1.

A preferred compound according to the present invention is a compound selected from the group consisting of:

3-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
4-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
4-{3-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
4-{4-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
4-(2-{4-[(3-Methyl-butylamino)-methyl]-phenoxy}-ethyl)-benzamide,
2-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
3-{3-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
3-(4-{[(Methyl-(3-methyl-butyl)-amino]-methyl}-phenoxy-methyl)-benzamide,
3-{4-[(3,3-Dimethyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
3-(4-{[2-(Tetrahydro-pyran-4-yl)-ethylamino]-methyl}-phenoxy-methyl)-benzamide,
3-(4-{[2-(4-Fluoro-phenyl)-ethylamino]-methyl}-phenoxy-methyl)-benzamide,
3-{3-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
3-{4-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
4-{3-[2-(3-Methyl-butylamino)-ethyl]-benzyloxy}-benzamide,
3-[4-(2-Benzylamino-ethyl)-Phenoxy-methyl]-benzamide, or a pharmaceutically acceptable salt, solvate, enantiomer, diastereomer and diastereomeric mixture thereof.


Preparing Compounds of the Invention

Compounds of formula I may be prepared as described in the following schemes and/or examples or following a combination of schemes known to one of skill in the art for making fragments and combinations thereof. Compounds employed as initial starting materials in the synthesis of compounds of the invention are well known and, to the extent not commercially available, are readily synthesized using specific references provided, or by standard procedures commonly employed by those of ordinary skill in the art and/or found in general reference texts.

More particularly, the compounds of the invention are produced in accordance with schemes 1 through 3 that are described in detail below, or analogous methods thereto. These reactions are often carried out following known procedures, methods, or analogous methods thereto. Examples of such known procedures and/or methods include those described in general reference texts such as Comprehensive Organic Transformations, VCH Publishers Inc, 1989; Compendium of Organic Synthetic Methods, Volumes 1-10, 1974-2002, Wiley Interscience; Advanced Organic Chemistry, Reactions Mechanisms, and Structure, 5th Edition, Michael B. Smith and Jerry March, Wiley Interscience, 2001; Advanced Organic Chemistry, 4th Edition, Part B, Reactions and Synthesis, Francis A. Carey and Richard J. Sundberg, Kluwer Academic / Plenum Publishers, 2000, etc., and references cited therein.


Compounds of formula I wherein k is equal 1 and j equal 0 may be prepared as shown for example in Scheme 1.

Scheme 1




Optionally substituted 4-hydroxy benzaldehyde **1** is reacted with optionally substituted hydroxymethyl benzonitrile **2**. The resulting nitrile compound **3** is then converted to the carboxamide **4** by hydrolysis procedures known to one of skill in the art, including for example, the use of hydrogen peroxide and potassium carbonate in a suitable solvent such as DMSO. The resulting amide compound **4** is reductively aminated at the aldehyde with an optionally substituted amine (primary and secondary) in a suitable solvent such as methanol and a reducing agent such as for example, sodium cyanoborohydride. The reductive amination of compound **4** results in the compound of formula **5**, which is a compound of the invention.

Compounds of formula I wherein j is 1 and k is 0 may be prepared as shown in scheme 2.



According to scheme 2, hydroxymethyl benzaldehydes (6) is coupled to an optionally substituted hydroxy-benzonitrile (7). The reaction is preferably effected by using Mitsunobu type reaction (triphenylphosphine and ligand (DEAD) in polar aprotic solvents such as THF to afford the coupled product 8. The nitrile group of 8 is hydrolyzed under basic conditions as described previously to afford the carboxamide 9. The carboxamide 9 is then reductively aminated to afford the desired amine 10, a compound of formula I. The starting material aldehyde 6 may be prepared by hydride reduction (DIBAL reduction) of the corresponding optionally substituted hydroxymethylbenzonitrile in toluene at temperatures of about -15 to 15°C .


Alternatively, compound 5 is utilized to afford a disubstituted compound of the invention as shown in Scheme 3 below.



As shown in Scheme 3, compound 5 upon reductive amination affords the amine 11 (wherein neither R^1 nor R^2 is hydrogen).

Compounds of formula I wherein k is greater than 1, may be prepared for example, according to scheme 4 below:


Scheme 4



As shown in scheme 4, the use of a substituted hydroxybenzaldehyde affords the ether compound 13 when reacted with a hydroxyethyl benzonitrile. The reaction is carried out using triphenylphosphine, DEAD and THF as solvent. The cyano group of the compound 13 is hydrolyzed under basic conditions using H_2O_2 and potassium carbonate, and DMSO as solvent to afford the carboxamide 14. Compound 14 is then reductively aminated with a suitable amine to afford the desired compound 15.

Certain compounds of the invention may also be accessed by protocols such as Scheme 5.


Scheme 5



For example, compounds of formula I having p greater than 1, may be more readily accessed by a Michael addition of nitromethane on an aldehyde 1 as shown in Scheme 5. The resulting nitroalkene 17 is reduced to the amine 18, and optionally protected for example by use of a Boc- group as the compound 19. The protected amino compound 19 is then reacted with an appropriately substituted hydroxymethylbenzonitrile to afford the intermediate 20 which is then deprotected at the Boc group to afford the compound 21. The compound 21 upon hydrolysis of the cyano group to the carboxamide 22 followed by reductive amination affords the desired product 23.

Compounds of formula I wherein k is 1 and j is 1 may be prepared as shown for example in the following Scheme 6.

Scheme 6



According to scheme 6, an optionally substituted hydroxymethyl benzaldehyde (6) is reacted with optionally substituted cyanobenzyl bromide (24) using cesium fluoride absorbed in Celite, and using acetonitrile as solvent. The reaction mixture is preferably heated to about reflux temperature to afford the coupled compound 25. The nitrile group of 25 is hydrolyzed under basic conditions as described previously to afford product 26. Compound 26 is then reductively aminated with a suitable amine to afford the desired compound 27.

Method of Using the Invention

As noted above, the compounds of the present invention are useful in blocking the effect of agonists at mu, kappa, and/or delta opioid receptors. As such, the present invention also provides a method for blocking a mu, kappa, delta receptor or receptor combination (heterodimer) thereof in a mammal comprising administering to said mammal a receptor blocking dose of a compound of formula I.

The term "receptor blocking dose", as used herein, means an amount of a compound of formula I necessary to block a mu, kappa, or delta receptor or receptor combination (heterodimer) thereof following administration to a mammal requiring blocking of a mu, kappa, or delta receptor or receptor combination (heterodimer) thereof.

The compounds of formula I or combinations thereof, are effective over a wide dosage range. For example, dosages per day will normally fall within the range of about 0.05 to about 250 mg/kg of body weight. In the treatment of adult humans, the range of about 0.5 to about 100 mg/kg, in single or divided doses, is preferred. However, it will be

understood that the amount of the compound actually administered will be determined by a physician in light of the relevant circumstances, including the condition to be treated, the choice of compound to be administered, the age, weight, response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration. Therefore, the above dosage ranges are not intended to limit the scope of the invention in any way. The compounds may be administered by a variety of routes such as the oral, transdermal, subcutaneous, intranasal, intramuscular and intravenous routes.

A variety of physiologic functions have been shown to be subject to or influenced by mu, kappa, or delta receptors or receptor combination (heterodimers) in the brain. As such, the compounds of the present invention are believed to have the ability to treat disorders associated with these receptors or combinations thereof such as eating disorders, opioid overdose, depression, smoking, alcoholism, sexual dysfunction, shock, stroke, spinal damage and head trauma. As such, the present invention also provides methods of treating the above disorders by blocking the effect of agonists at a mu, kappa, delta receptors or receptor combinations (heterodimer) thereof.

Assay Methodology

The compounds of the present invention have been found to display significant activity in an opioid receptor binding assay which measures the ability of the compounds to block the mu, kappa, delta or receptor combination (heterodimer) thereof. The functional antagonist potency (Kb) of the sample compounds was determined using the GTP γ S binding assay. GTP γ S – based functional assays provide an in vitro measure of the activity of opioid agonists and antagonists. Opioid reference compounds or test compound are incubated with membrane homogenate from cells expressing the cloned human mu, kappa or delta opioid receptor and radiolabeled [35S]GTP γ S. If the compound activates the opioid receptor, an increase in the binding of radiolabeled GTP γ S is observed. Similarly, if the compound exhibits antagonist activity, it interferes with the ability of the control agonist to stimulate GTP γ S binding. These tests provide an in vitro measurement of the activity of the compound at human opioid receptors.

GTP- γ -S Binding Assay

An SPA - based GTP- γ -S assay format was developed based on previous opioid (Emmerson et al., J. Pharm Exp Ther 278,1121,1996; Horng et al., Society for Neuroscience Abstracts, 434.6, 2000) and muscarinic (DeLapp et al., JPET 289, 946, 1999) assay formats. Membranes were resuspended in 20 mM HEPES, 100 mM NaCl, 5 mM MgCl₂, 1 mM DTT, and 1 mM EDTA. Fifty (50) μ L of GTP- γ -[35S], compound, membrane suspension (20 microgram/well), and wheat germ agglutinin coated SPA beads (1mg/well) were added to clear bottom 96 well assay plates. GDP (200 mM) was added to the membrane solution prior to addition to the assay plates. Plates were sealed and incubated for four hours at room temperature then placed in a refrigerator overnight to allow the beads to settle. Signal stability at 4 °C was determined to be > 60 hours. Plates were warmed to room temperature and counted in a Wallac Microbeta scintillation counter. For antagonist assays, specific agonists were added at the following concentrations: (MOR) DAMGO 1 micromolar, (DOR) DPDPE 30 nM, (KOR) U69593 300 nM. K_b's were determined by Cheng-Prusoff equation (see Cheng and Prusoff, Biochem. Pharmacol. 22, 3099, 1973). Results obtained for a sample of compounds of the invention in the GTP- γ -S Binding Assay are shown in table 1 below.

Table 1

Example #	K _b (nM) mu	K _b (nM) kappa	K _b (nM) delta
1	>21		>41
2	62.86		336.65
3	13.56	14.04	194.00
4	>23		>67
5	>23		>67

Formulation

A compound of the invention is preferably presented in the form of a pharmaceutical formulation comprising a pharmaceutically acceptable carrier, diluent or excipient and a compound of the invention. Such compositions will contain from about

0.1 percent by weight to about 90.0 percent by weight of the compound of the invention (Active Ingredient). As such, the present invention also provides pharmaceutical formulations comprising a compound of the invention and a pharmaceutically acceptable carrier, diluent or excipient therefore.

In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material that acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, emulsions, solutions, syrups, suspensions, aerosols (as a solid or in a liquid medium), and soft and hard gelatin capsules.

Examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, tragacanth, gelatin, syrup, methyl cellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate, water, and mineral oil. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The formulations of the invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

For oral administration, the Active Ingredient, a compound of this invention, may be admixed with carriers and diluents and molded into tablets or enclosed in gelatin capsules.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 1 to about 500 mg, more usually about 5 to about 300 mg, of the Active Ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier.

In order to more fully illustrate the operation of this invention, the following formulation examples are provided. The examples are illustrative only, and are not

intended to limit the scope of the invention. The formulations may employ as Active Ingredient any of the compounds of the present invention.

FORMULATION 1

Hard gelatin capsules are prepared using the following ingredients:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	250	55
Starch dried	200	43
Magnesium stearate	10	2

The above ingredients are mixed and filled into hard gelatin capsules in 460 mg quantities.

FORMULATION 2

Capsules each containing 20 mg of medicament are made as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	20	10
Starch	89	44.5
Microcrystalline cellulose	89	44.5
Magnesium stearate	2	1

The active ingredient, cellulose, starch and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve and filled into a hard gelatin capsule.

FORMULATION 3

Capsules each containing 100 mg of active ingredient are made as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	100	30
Polyoxyethylene Sorbitan monooleate	50mcg	0.02
Starch powder	250	69.98

The above ingredients are thoroughly mixed and placed in an empty gelatin capsule.

FORMULATION 4

Tablets each containing 10 mg of active ingredient are prepared as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	10	10
Starch	45	45
Microcrystalline cellulose	35	35
Polyvinylpyrrolidone (as 10% solution in water)	4	4
Sodium carboxymethyl starch	4.5	4.5
Magnesium stearate	0.5	0.5
talc	1	1

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders, which are then passed through a No. 14 mesh U.S. sieve. The granule so produced is dried at 50-60°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules, which after mixing, is compressed on a tablet machine to yield a tablet weighing 100 mg.

FORMULATION 5

A tablet formula may be prepared using the ingredients below:

Compound	Amount per capsule (mg)	Percent by weight (%)
Active Ingredient	250	38
Cellulose microcrystalline	400	60
Silicon dioxide fumed	10	1.5
Stearic acid	5	0.5

The components are blended and compressed to form tablets each weighing 665mg.

FORMULATION 6

Suspensions each containing 5 mg of medicament per 5 ml dose are made as follows:

Compound	Amount per 5mL suspension (ml)
Active Ingredient	5
Sodium carboxymethyl cellulose	50
Syrup	1.25
Benzoic acid solution	0.10
Flavor	q.v.
Color	q.v.
Water	q.s. to 5mL

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethylcellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color is diluted with some of the water and added to the paste with stirring. Sufficient water is then added to produce the required volume.

FORMULATION 7

An aerosol solution is prepared containing the following components:

Compound	Concentration by weight
----------	-------------------------


	(percent)
Active Ingredient	0.25
Ethanol	29.75
Propellant 22 (chlorodifluoromethane)	70.0

The active compound is mixed with ethanol and the mixture added to a portion of the Propellant 22, cooled to -30°C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted further with the remaining amount of propellant. The valve units are then fitted to the container.

Examples


Example 1

3-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide



Step 1

3-(4-Formyl-phenoxy-methyl)-benzonitrile




To a solution of 3-(Hydroxymethyl)benzonitrile (1.1 equiv), triphenylphosphine (1.1 equiv), and 4-Hydroxybenzaldehyde (1 equiv) in dry THF was added dropwise Diethyl azodicarboxylate (DEAD) (1.4 equiv). The reaction mixture stirred at room temperature overnight. The reaction was monitored by TLC. The solvent was removed under vacuum. It was purified by silica gel chromatography using Hexane: Ethyl acetate (3:1) to afford the title compound as a white solid.

58 % Yield

^1H NMR (CDCl_3 , 300 MHz) δ : 9.91 (s, 1H), 7.87 (d, $J = 9.0$ Hz, 2H), 7.69 (s, 1H), 7.68-7.63 (m, 2H), 7.54 (d, $J = 7.8$ Hz, 1H), 7.07 (d, $J = 8.7$ Hz, 2H), 5.17 (s, 2H).

Step 2

3-(4-Formyl-phenoxy-methyl)-benzamide




To a solution of 3-(4-Formyl-phenoxy-methyl)benzonitrile (1 equiv), in DMSO was added potassium carbonate (0.5 equiv), and hydrogen peroxide (4 equiv). The reaction mixture stirred at room temperature for 2 hours. The reaction was monitored by TLC. It was poured on water and extracted with ethyl acetate (x 3). The organic phase was dried over magnesium sulphate, filtered and concentrated under vacuum to yield a white solid.

99 % Yield

^1H NMR (CDCl_3 , 300 MHz) δ : 9.84 (s, 1H), 7.99-7.98 (m, 1H), 7.88 (d, $J = 9.0$ Hz, 2H), 7.86-7.83 (m, 1H), 7.64 (m, 1H), 7.50 (m, 1H), 7.18 (d, $J = 8.4$ Hz, 2H), 5.17 (s, 2H).

MS (Electrospray): ($\text{M}^+ - 1$) 254.3

Step 3



A mixture of 3-(4-formyl-phenoxy-methyl)-benzamide (1 equiv), isoamylamine (1 equiv), 4 Å molecular sieves (1000% weight) in methanol (0.1 M) was stirred overnight under nitrogen atmosphere at room temperature. The following day NaBH_4 (5 equiv) was added and the reaction mixture was stirred for 3 hours. The reaction can be monitored by TLC analysis. The reaction mixture was filtered off and the solvent evaporated to yield a residue which was purified by silica gel chromatography using CHCl_3 : EtOH 7%: NH_4OH 0.7 to afford the title compound as a solid.


75% Yield

^1H NMR (CD_3OD , 300 MHz) δ : 7.96 (s, 1H), 7.82-7.80 (m, 1H), 7.64-7.61 (m, 1H), 7.49-7.46 (m, 1H), 7.25 (d, $J = 8.7$ Hz, 2H), 6.97 (d, $J = 9.0$ Hz, 2H), 5.13 (s, 2H), 3.69 (s, 2H), 2.61-2.56 (m, 2H), 1.61-1.54 (m, 1H), 1.45-1.40 (m, 2H), 0.89 (d, $J = 6.6$ Hz, 6H).

MS (Electrospray): ($\text{M}^+ + 1$) 327.3

Example 2

4-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide



The compound of Example 2 may be prepared essentially as described in Example

1.

42% Yield


^1H NMR (CD_3OD , 300 MHz) δ : 7.87 (d, $J = 8.4$ Hz, 2H), 7.52 (d, $J = 8.7$ Hz, 2H), 7.24 (d, $J = 8.7$ Hz, 2H), 6.95 (d, $J = 8.7$ Hz, 2H), 5.14 (s, 2H), 3.66 (s, 2H), 2.58-2.53 (m, 2H), 1.58-1.56 (m, 1H), 1.43-1.36 (m, 2H), 0.88 (d, $J = 6.6$ Hz, 6H).

MS (Electrospray): ($\text{M}^+ + 1$) 327.3

HPLC = 98% @ 5.95m (5/95 to 95/5 ACN/(0.1%TFA in water) over 10 minutes, Zorbax SB-Phenyl 4.6mmx15cmx5micron, $\lambda=254\text{nm}$.de

Example 3

4-{3-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide



The compound of Example 3 may be prepared essentially as described in Example

1.

32% Yield


$^1\text{H NMR}$ (CD_3OD , 300 MHz) δ : 7.83 (d, $J = 8.7$ Hz, 2H), 7.43 (s, 1H), 7.36-7.34(m, 2H), 7.34-7.30 (m, 1H), 7.04 (d, $J = 9.0$ Hz, 2H), 5.15 (s, 2H), 3.76 (s, 2H), 2.59-2.56 (m, 2H), 1.60-1.55 (m, 1H), 1.44-1.40 (m, 2H), 0.92 (d, $J = 6.6$ Hz, 6H).

MS (Electrospray): ($\text{M}^+ + 1$) 327.2

HPLC = 94% @ 5.96m (5/95 to 95/5 ACN/(0.1%TFA in water) over 10 minutes, Zorbax SB-Phenyl 4.6mmx15cmx5micron, $\lambda=254\text{nm}$.de

Example 4

4-{4-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide



The compound of Example 4 may be prepared essentially as described in Example

1.

59% Yield


^1H NMR (CD_3OD , 300 MHz) δ : 7.85 (d, J = 8.4 Hz, 2H), 7.42-7.38 (m, 4H), 7.04 (d, J = 8.7 Hz, 2H), 5.15 (s, 2H), 3.79 (s, 2H), 2.58-2.53 (m, 2H), 1.65-1.58 (m, 1H), 1.46-1.40 (m, 2H), 0.90 (d, J = 6.6 Hz, 6H).

MS (Electrospray): (M^+ +1) 327.2

HPLC = 97% @ 6.02m (5/95 to 95/5 ACN/(0.1%TFA in water) over 10 minutes, Zorbax SB-Phenyl 4.6mmx15cmx5micron, λ =254nm.de

Example 5

4-(2-{4-[(3-Methyl-butylamino)-methyl]-phenoxy}-ethyl)-benzamide



The compound of Example 5 may be prepared essentially as described in Example 1.

71% Yield


^1H NMR (CD_3OD , 300 MHz) δ : 7.81 (d, J = 8.1 Hz, 2H), 7.40 (d, J = 8.1 Hz, 2H), 7.23 (d, J = 8.7 Hz, 2H), 6.87 (d, J = 8.7 Hz, 2H), 5.15 (s, 2H), 4.21 (t, J = 6.6 Hz, 2H), 3.70 (s, 2H), 3.12 (t, J = 6.6 Hz, 2H), 2.63-2.58 (m, 2H), 1.65-1.58 (m, 1H), 1.45-1.40 (m, 2H), 0.89 (d, J = 6.6 Hz, 6H).

MS (Electrospray): (M^+ +1) 341.3 (M^+ -1) 339.4

HPLC = 91% @ 6.07m (5/95 to 95/5 ACN/(0.1%TFA in water) over 10 minutes, Zorbax SB-Phenyl 4.6mmx15cmx5micron, λ =254nm.de

We claim:

1. A compound of formula (I)



p is 0, 1, or 2;

y is 0, 1, or 2; and z is 0, 1, or 2;

j and k are each independently 0, 1, or 2 provided that j and k are not simultaneously equal to 0;

X₁ and X₂ are each independently is CH, or N;

R¹ and R² are independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, C₁-C₁₀ alkylaryl, SO₂R⁸, (CH₂)_nC(O)NR⁸R⁸, SO₂C₁-C₁₀ alkylaryl, SO₂C₁-C₈ alkylheterocyclic, C₄-C₁₀ alkylcycloalkane, (CH₂)_nC(O)OR⁸, and (CH₂)_nC(O)R⁸; wherein each of the alkyl, alkenyl, and aryl groups are optionally

substituted with one to two groups independently selected from C₁-C₈ alkyl, C₂-C₈ alkenyl, phenyl, C₃-C₈ cycloalkyl, C₁-C₈ alkylaryl, and C(O)C₁-C₈ alkyl; and wherein R¹ and R² may optionally combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may further have substituents selected from the group consisting of oxo, amino, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, C₁-C₃ alkylaryl, C(O)C₁-C₈ alkyl, CO(O)C₁-C₈ alkyl, halo, C₁-C₃ haloalkyl;

R³ and R^{3'} are each independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, aryl, C₁-C₈ alkylcycloalkyl, and C₁-C₈ alkylaryl;


R⁴ and R⁵ are each independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₁-C₈ alkoxy, halo, C₁-C₈ haloalkyl, phenyl, aryl, C₁-C₈ alkylaryl, (CH₂)_mNSO₂C₁-C₈ alkyl, (CH₂)_mNSO₂phenyl, (CH₂)_mNSO₂aryl, -C(O)C₁-C₈ alkyl, and -

$C(O)OC_1-C_8$ alkyl; wherein each R^4 and R^5 is attached to its respective ring only at carbon atoms; wherein m is 1 or 2; and n is 0, 1, 2, or 3;

R^6 and R^7 are each independently selected from hydrogen, C_1-C_8 alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, $C(O)C_1-C_8$ alkyl, $SO_2C_1-C_8$ alkyl, $SO_2C_1-C_8$ alkylaryl, $SO_2C_1-C_8$ alkylheterocyclic, aryl, C_1-C_8 alkylaryl, C_3-C_7 cycloalkane, C_1-C_6 alkylcycloalkane, $(CH_2)_mC(O)OR^8$, $(CH_2)_mC(O)R^8$, $(CH_2)_mC(O)NR^8R^8$, and $(CH_2)_mNSO_2R^8$; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to two groups independently selected from C_1-C_8 alkyl, C_2-C_8 alkenyl, phenyl, and C_1-C_8 alkylaryl; and wherein R^6 and R^7 may independently combine with each other, and with the nitrogen atom to which they are attached to form a 4, 5, 6, or 7-membered nitrogen containing heterocycle which nitrogen containing heterocycle may optionally have substituents selected from the group consisting of oxo, amino, C_1-C_8 alkyl, C_2-C_8 alkenyl, C_2-C_8 alkynyl, phenyl, and C_1-C_8 alkylaryl;

R^8 is independently selected from hydrogen, C_1-C_8 alkyl, C_2-C_8 alkenyl, phenyl, benzyl, and C_5-C_8 alkylaryl; or a pharmaceutically acceptable salt, solvate, prodrug, enantiomer, racemate, diastereomer, or mixture of diastereomers thereof.

2. The compound according to claim 1 wherein the X_1 is selected CH.
3. A compound according to Claim 1 wherein X_1 is N.
4. A compound according to Claim 1 wherein y is 0 or 1, and R^4 is independently selected from the group consisting of fluoro, chloro, bromo, methoxy, ethoxy, methyl, ethyl, isopropyl, trifluoromethyl, phenyl, benzyl and ethoxy.
5. A compound according to Claim 1 wherein z is 0 or 1, and R^5 is independently selected from the group consisting of fluoro, chloro, bromo, methoxy, ethoxy, methyl, ethyl, isopropyl, trifluoromethyl, phenyl, and benzyl.
6. A compound according to Claim 1 wherein R^1 and R^2 are each independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, 2-methylpentyl, t-butyl, cyclopropyl, phenyl,



7. The compound according to Claim 1 wherein R^6 and R^7 are each independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, and phenyl.

8. A compound selected from the group consisting of:

- 3-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
- 4-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
- 4-{3-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
- 4-{4-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
- 4-(2-{4-[(3-Methyl-butylamino)-methyl]-phenoxy}-ethyl)-benzamide,
- 2-{4-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
- 3-{3-[(3-Methyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
- 3-(4-{[(Methyl-(3-methyl-butyl)-amino]-methyl}-phenoxy-methyl)-benzamide,
- 3-{4-[(3,3-Dimethyl-butylamino)-methyl]-phenoxy-methyl}-benzamide,
- 3-(4-{[2-(Tetrahydro-pyran-4-yl)-ethylamino]-methyl}-phenoxy-methyl)-benzamide,
- 3-(4-{[2-(4-Fluoro-phenyl)-ethylamino]-methyl}-phenoxy-methyl)-benzamide,
- 3-{3-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
- 3-{4-[(3-Methyl-butylamino)-methyl]-benzyloxy}-benzamide,
- 4-{3-[2-(3-Methyl-butylamino)-ethyl]-benzyloxy}-benzamide,

3-[4-(2-Benzylamino-ethyl)-Phenoxy-methyl]-benzamide, or a pharmaceutically acceptable salt, solvate, enantiomer, diastereomer and diastereomeric mixture thereof.

9. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Claim 1 in association with a carrier, diluent and/or excipient.

10. A method for blocking a mu, kappa, delta or receptor combination (heterodimer) thereof in mammals comprising administering to a mammal requiring blocking of a mu, kappa, delta or receptor combination (heterodimer) thereof, a receptor blocking dose of a compound of formula I, or a pharmaceutically acceptable salt, enantiomer, racemate, mixture of diastereomers, or solvate thereof.

11. A method of treating or preventing obesity and Related Diseases comprising administering a therapeutically effective amount of a compound of Claim 1.

12. A method according to Claim 11 wherein the Related Diseases is selected from the group consisting of diabetes, diabetic complications, diabetic retinopathy, atherosclerosis, hyperlipidemia, hypertriglycemia, hyperglycemia, and hyperlipoproteinemia.

13. A method of treating and/or preventing diseases related to obesity including irritable bowel syndrome, nausea, vomiting, depression, smoking and alcohol addiction, sexual dysfunction, substance abuse, drug overdose, addictive behavior disorders, compulsive behaviors, and stroke comprising administering a therapeutically effective amount of a compound of formula I.

14. A method of suppressing appetite in a patient in need thereof, comprising administering a therapeutically effective amount of a compound of formula I.

15. Use of a compound of formula I in the manufacture of a medicament for the treatment and/or amelioration of the symptoms associated with obesity and Related

Diseases comprising administering a therapeutically effective amount of a compound of formula I to a patient in need thereof.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C235/44 C07C235/46 A61K31/166 A61P3/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, CHEM ABS Data, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE BEILSTEIN BEILSTEIN INSTITUT ZUR FÖRDERUNG DER CHEMISCHEN WISSENSCHAFT, FRANKFURT AM MAIN, DE; XP002334306 retrieved from XFIRE Database accession no. 5541857, 5745449 abstract & J. MED. CHEM., vol. 27, no. 2, 1984, pages 129-143,</p>	1, 2, 4-7, 9
X	<p>GB 1 504 604 A (CIBA-GEIGY AG) 22 March 1978 (1978-03-22) Compound 7 in example 10</p> <p style="text-align: center;">-/--</p>	1, 2, 4-7, 9-14

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.*** Special categories of cited documents:**

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

4 July 2005

Date of mailing of the international search report

10/08/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Cooper, S

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 965 741 A (BREAULT ET AL) 12 October 1999 (1999-10-12) Table VI, compound 17; column 18, line 35 - column 21, line 8; claim 1	1-7,9-14
P,A	----- WO 2004/026305 A (ELI LILLY AND COMPANY; BLANCO-PILLADO, MARIA-JESUS; CHAPPELL, MARK, DO) 1 April 2004 (2004-04-01) abstract -----	1-15

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 10-14 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1504604	A	22-03-1978	AT 340388 B 12-12-1977
			AT 604275 A 15-04-1977
			AT 340390 B 12-12-1977
			AT 783276 A 15-04-1977
			AT 340391 B 12-12-1977
			AT 783376 A 15-04-1977
			AT 340392 B 12-12-1977
			AT 783576 A 15-04-1977
			AT 340393 B 12-12-1977
			AT 783676 A 15-04-1977
			AU 8362675 A 10-02-1977
			BE 832082 A1 04-02-1976
			CA 1069923 A1 15-01-1980
			CH 612910 A5 31-08-1979
			CH 612911 A5 31-08-1979
			CH 612912 A5 31-08-1979
			CH 612913 A5 31-08-1979
			CH 612909 A5 31-08-1979
			DE 2534339 A1 19-02-1976
			DK 317175 A 06-02-1976
			ES 439997 A1 16-06-1977
			ES 455488 A1 01-01-1978
			ES 455489 A1 01-01-1978
			ES 455490 A1 01-01-1978
			ES 455491 A1 01-01-1978
			FI 752095 A 06-02-1976
			FR 2281105 A1 05-03-1976
			HU 175581 B 28-09-1980
			IE 42335 B1 16-07-1980
			IL 47813 A 25-07-1979
			JP 51041345 A 07-04-1976
			NL 7509270 A 09-02-1976
			NO 752621 A 06-02-1976
			SE 7508181 A 06-02-1976
			US 4329367 A 11-05-1982
			ZA 7504192 A 30-06-1976
			ZA 7505018 A 28-07-1976
US 5965741	A	12-10-1999	AT 185791 T 15-11-1999
			AU 3351995 A 22-03-1996
			DE 69512925 D1 25-11-1999
			DE 69512925 T2 04-05-2000
			EP 0778821 A1 18-06-1997
			WO 9606822 A1 07-03-1996
			JP 10504836 T 12-05-1998
WO 2004026305	A	01-04-2004	AU 2003269980 A1 08-04-2004
			CA 2499690 A1 01-04-2004
			WO 2004026305 A1 01-04-2004